

# Your Birth Control Choices

Method	How well does it work?	How to Use	Pros	Cons
<b>The Implant</b> Nexplanon® 	> 99%	A health care provider places it under the skin of the upper arm It must be removed by a health care provider	Long lasting (up to 5 years) No pill to take daily Often decreases cramps Can be used while breastfeeding You can become pregnant right after it is removed	Can cause irregular bleeding After 1 year, you may have no period at all Does not protect against human immunodeficiency virus (HIV) or other sexually transmitted infections (STIs)
<b>Progestin IUD</b> Liletta®, Mirena®, Skyla® and others  <p>Mirena</p>	> 99%	Must be placed in uterus by a health care provider Usually removed by a health care provider	May be left in place 3 to 7 years, depending on which IUD you choose No pill to take daily May improve period cramps and bleeding Can be used while breastfeeding You can become pregnant right after it is removed	May cause lighter periods, spotting, or no period at all Rarely, uterus is injured during placement Does not protect against HIV or other STIs
<b>Copper IUD</b> ParaGard® 	> 99%	Must be placed in uterus by a health care provider Usually removed by a health care provider	May be left in place for up to 12 years No pill to take daily Can be used while breastfeeding You can become pregnant right after it is removed	May cause more cramps and heavier periods May cause spotting between periods Rarely, uterus is injured during placement Does not protect against HIV or other STIs
<b>The Shot</b> Depo-Provera® 	94%	Get a shot every 3 months	Each shot works for 12 weeks Private Usually decreases periods Helps prevent cancer of the uterus No pill to take daily Can be used while breastfeeding	May cause spotting, no period, weight gain, depression, hair or skin changes, change in sex drive May cause delay in getting pregnant after you stop the shots Side effects may last up to 6 months after you stop the shots Does not protect against HIV or other STIs
<b>The Pill</b> 	91%	Must take the pill daily	Can make periods more regular and less painful Can improve PMS symptoms Can improve acne Helps prevent cancer of the ovaries You can become pregnant right after stopping the pills	May cause nausea, weight gain, headaches, change in sex drive – some of these can be relieved by changing to a new brand May cause spotting the first 1-2 months Does not protect against HIV or other STIs
<b>Progestin-Only Pills</b> 	91%	Must take the pill daily	Can be used while breastfeeding You can become pregnant right after stopping the pills	Often causes spotting, which may last for many months May cause depression, hair or skin changes, change in sex drive Does not protect against HIV or other STIs
<b>The Patch</b> Ortho Evra® 	91%	Apply a new patch once a week for three weeks No patch in week 4	Can make periods more regular and less painful No pill to take daily You can become pregnant right after stopping patch	Can irritate skin under the patch May cause spotting the first 1-2 months Does not protect against HIV or other STIs
<b>The Ring</b> Nuvaring® 	91%	Insert a small ring into the vagina Change ring each month	One size fits all Private Does not require spermicide Can make periods more regular and less painful No pill to take daily You can become pregnant right after stopping the ring	Can increase vaginal discharge May cause spotting the first 1-2 months of use Does not protect against HIV or other STIs